

# BULLETIN OF THE OKLAHOMA ORNITHOLOGICAL SOCIETY

VOL. 34

DECEMBER 2001

No. 4

*Bull. Okla. Ornithol. Soc.* 34:21–25, 2001  
© Oklahoma Ornithological Society 2001

## Recent documentation restores the Broad-tailed Hummingbird to Oklahoma's state bird list

BY DAN L. REINKING, JOHN E. PUSCHOCK, AND BONNIE L. GALL

On 3 September 1999, while birding in the Black Mesa area of Cimarron County in the western Oklahoma Panhandle, we stopped at the Kenton Mercantile around midday to observe the hummingbird feeders. We positioned ourselves to obtain good views of the feeders and began to watch approximately four to six Black-chinned Hummingbirds (*Archilochus alexandri*) that were actively seeking syrup. After some time, Puschock called our attention to a different hummingbird that had appeared. We each observed the bird with binoculars from the parking area in front of the store from a distance of about 10 m away with the sun overhead and providing good light even though the feeders were shaded under the porch of the store. The bird was observed perched on the feeder as well as hovering, and it flared its tail several times with its back towards us while hovering as it tried to displace a wasp from the feeder. The total observation time was about one minute, but we had close, unobstructed views in good light for the whole duration. The bird then flew away and did not return while we waited. A cold front came through that night. When we visited the store the following day, two Black-chinned Hummingbirds were the only hummingbirds we saw.

Immediately after our observation of the unusual hummingbird, we independently recorded descriptions of the bird, and only then compared notes and reviewed field guides. All observers noted green upperparts, including the crown, nape, back, and rump. Puschock and Gall each noted that the throat was off-white with a few dark feathers. Reinking described a broad, pale cinnamon wash on the flanks. Puschock described the flanks as the "diffuse burnt orange/rufous" color typical for *Selasphorus* hummingbirds, although less intense in tone than that seen on adult female Rufous Hummingbirds (*S. rufus*). He also noted that the demarcation of the flank color was less sharp than on adult female Rufous Hummingbirds. Gall described the flanks as buffy.

During the observation, Puschock suggested that we look carefully at the tail, and all observers independently noted white-tipped outer tail feathers with some rufous (less than half of the tail length) on the proxi-

mal portion of the tail. Puschock specifically noted three outer tail feathers with rufous. All observers described the central tail feathers as either green or "dark" in color, lacking either rufous or white. The tail features were seen well by all three observers, as the bird flared its tail several times while facing directly away from us. Reinking noted the relatively broad width of the outer tail feathers, a feature he had previously evaluated in the hand on Allen's (*S. sasin*) and Rufous hummingbirds while banding in California. All observers also independently noted that the bird appeared larger than the Black-chinned Hummingbirds which were present at the time of the observation. Reinking specifically noted the overall longer body length as compared to the Black-chinned Hummingbirds.

Several key features, including the presence of obvious rufous coloration on the flanks, the large size relative to Black-chinned Hummingbirds, the dark rufous in the tail being restricted to relatively small, proximal portions of three outer tail feathers, and the lack of any rufous in the back or rump all led us to conclude that this was in fact a Broad-tailed Hummingbird (*S. platycercus*). Immature *Archilochus* hummingbirds may occasionally show buffy coloration on the flanks but usually not to the extent and intensity of *Selasphorus* hummingbirds (Kaufman 1990). *Archilochus* hummingbirds may rarely show some buffy or pale rufous tones in the tail, but not the more extensive, dark chestnut rufous of *Selasphorus* hummingbirds such as was observed in the tail of this bird. Both Allen's and Rufous hummingbirds would have more extensive rufous in the tail, often including some in the inner tail feathers, would show more intense rufous coloration on the flanks, and usually would show at least some rufous on the back or uppertail coverts (Pyle 1997; Sibley 2000; Williamson in press). Calliope Hummingbirds (*Stellula caliope*) have very little rufous in the tail (Williamson 1999; Sibley 2000), less than that observed on this bird. Finally, Broad-tailed Hummingbirds are larger than Calliope, Allen's, or Rufous hummingbirds (and *Archilochus* hummingbirds), due in part to a longer tail (Howell and Webb 1995; Pyle 1997; Sibley 2000; Williamson in press). While the tail length of this bird was not specifically described, two observers made generic note of the larger overall size, and one observer did specifically note a longer overall body length on this bird in relation to the Black-chinned Hummingbirds it was directly and closely compared with. This size difference alone is significant in helping eliminate other *Selasphorus* and *Archilochus* hummingbirds as possibilities. The fact that all three observers independently noted these multiple, important characteristics with great consistency is also significant in evaluating this record. The documentations submitted by the authors were subsequently reviewed and accepted by the Oklahoma Bird Records Committee (OBRC).

The Broad-tailed Hummingbird breeds in open woodlands and shrublands in the mountains of western North America and Mexico. In the parts of its range closest to Oklahoma, it nests in the mountains of

south central Colorado and northeastern New Mexico (Bailey 1928; Calder and Calder 1992; Boyle 1998). It is a rare but regular migrant in eastern New Mexico (J. Oldenettel and C. Rustay, pers. comm.). There are a number of substantiated Broad-tailed Hummingbird records in Kansas (Thompson and Ely 1989; Grzybowski 1989, 1992a,b). Seyffert (2001) considers the Broad-tailed Hummingbird an uncommon to fairly common migrant in the Texas Panhandle, having been seen in six counties. The American Ornithologists' Union (1998) describes the Broad-tailed Hummingbird as a casual migrant east to Nebraska, central Kansas, and eastern and southeastern Texas. Extralimital sightings to the east have also occurred in Alabama, Arkansas, Delaware, Georgia, Louisiana, Illinois, and Mississippi (Williamson in press; E. Womack, pers. comm.), and one specimen and one banding record during the year 2000 exist for Florida (G. Wallace and L. Duncan, pers. comm.). Williamson (in press) illustrates the eastern extent of most Broad-tailed Hummingbird migrants as passing through the eastern Panhandle of Oklahoma, presumably based on multiple records from Kansas and the Texas Panhandle.

In Oklahoma, Tate (1923) reported a nest with two eggs in Cimarron County in 1912 and a September sighting in Kenton in 1922. No supporting description of either sighting was provided. Based on the information from Tate as well as a specimen reportedly collected by Tate, Nice (1931) considered the Broad-tailed Hummingbird a "rare transient and summer resident in Cimarron County." The specimen has since been destroyed in a fire (Grzybowski et al. 1992). A 1932 hummingbird observation near Kenton by Sutton (1934) is later described by him as "probably of this species," and he also reported a 1953 sighting in Texas County (Sutton 1967). Baumgartner and Baumgartner (1992) do not provide any additional records of this species in Oklahoma.

In evaluating all reported Oklahoma records of the Broad-tailed Hummingbird for Grzybowski et al. (1992), the OBRC voted to delete this species from the Oklahoma bird list based on the absence of an existing specimen and the lack of supporting documentation for any of the reported sightings. The acceptance of this current record by the OBRC therefore restores the Broad-tailed Hummingbird to the Oklahoma state bird list in "hypothetical" status, meaning that the written documentations were convincing to at least a majority of committee members, although a specimen or identifiable photograph from Oklahoma does not yet exist (Oklahoma Bird Records Committee 2000).

Given the proximity of the Broad-tailed Hummingbird's breeding range to Oklahoma and the number of records reported from Kansas and western Texas, it is possible if not likely that this species is more common in the western Oklahoma Panhandle, at least during migration, than the existing records would indicate. The current availability of hummingbird feeders in the Kenton area provides an opportunity for careful observers to test this idea.

We thank Allan Griggs for placing hummingbird feeders at "The Merc." Comments from James Hoffman, Charles R. Brown, and Ellie Womack helped improve the manuscript. Sheri Williamson provided helpful information.

#### LITERATURE CITED

- American Ornithologists' Union. 1998. Check-List of North American birds. 7th ed. American Ornithologists' Union, Washington, D.C.
- Bailey, F. M. 1928. Birds of New Mexico. Judd and Detweiler, Washington, D.C.
- Baumgartner, F. M., and A. M. Baumgartner. 1992. Oklahoma bird life. Univ. Oklahoma Press, Norman.
- Boyle, S. 1998. Broad-tailed Hummingbird. *In* Colorado breeding bird atlas (H. E. Kingery, ed.). Colorado Bird Atlas Partnership, Denver.
- Calder, W. A., and L. L. Calder. 1992. Broad-tailed Hummingbird. *In* The birds of North America, No. 16 (A. Poole and F. Gill, eds.). Academy of Natural Sciences, Philadelphia, and the American Ornithologists' Union, Washington, D.C.
- Grzybowski, J. A. 1989. Southern Great Plains region. *Am. Birds* 43:124–126.
- Grzybowski, J. A. 1992a. Southern Great Plains region. *Am. Birds* 46:113–117.
- Grzybowski, J. A. 1992b. Southern Great Plains region. *Am. Birds* 46:1151–1152.
- Grzybowski, J. A., J. W. Arterburn, W. A. Carter, J. S. Tomer, and D. W. Verser. 1992. Date guide to the occurrences of birds in Oklahoma. 2nd ed. Oklahoma Ornithol. Soc., Norman.
- Howell, S. N. G., and S. Webb. 1995. A guide to the birds of Mexico and northern Central America. Oxford Univ. Press, New York.
- Kaufman, K. 1990. A field guide to advanced birding. Houghton Mifflin, Boston.
- Nice, M. M. 1931. The birds of Oklahoma. Revised Edition. *Publ. Oklahoma Biol. Surv.* 3(1).
- Oklahoma Bird Records Committee. 2000. Date guide to the occurrences of birds in Oklahoma. 3rd ed. Oklahoma Ornithol. Soc., Tulsa.
- Pyle, P. 1997. Identification guide to North American birds. Part I. Slate Creek Press, Bolinas, California.
- Seyffert, K. D. 2001. Birds of the Texas Panhandle: their status, distribution, and history. Texas A&M Univ. Press, College Station, Texas.
- Sibley, D. A. 2000. National Audubon Society Sibley guide to birds. Knopf, New York.
- Sutton, G. M. 1934. Notes on the birds of the western Panhandle of Oklahoma. *Ann. Carnegie Mus.* 24:1–50.
- Sutton, G. M. 1967. Oklahoma birds. Univ. Oklahoma Press, Norman.
- Tate, R. C. 1923. Some birds of the Oklahoma Panhandle. *Proc. Oklahoma Acad. Sci.* 3:41–51.

- Thompson, M. C., and C. Ely. 1989. *Birds in Kansas*. Vol. 1. Univ. Press of Kansas, Lawrence.
- Williamson, S. 1999. The feminine mystique: identifying female-plumaged hummingbirds. *WildBird*, May, 30–33.
- Williamson, S. in press. *A field guide to hummingbirds of North America*. Houghton Mifflin, New York.

GEORGE M. SUTTON AVIAN RESEARCH CENTER, P.O. BOX 2007, BARTLESVILLE, OK. 74005 (DLR); USDA FOREST SERVICE, SEMINOLE RANGER DISTRICT, 40929 STATE ROAD 19, UMATILLA, FL. 32784 (JEP); 403140 W. 2010 DR., BARTLESVILLE, OK. 74006 (BLG). Received 6 March 2001, accepted 17 October 2001.

## NOTES

*Bull. Okla. Ornithol. Soc.* 34:25–27, 2001  
© Oklahoma Ornithological Society 2001

**Wood Stork invasion in McCurtain County, Oklahoma.**—An invasion of Wood Storks (*Mycteria americana*) into McCurtain County, Oklahoma, occurred during the summer of 2000. The first storks reported that year were four that we observed at the Red Slough Wetland Reserve Project (Red Slough) on 5 June. By August, numbers had increased dramatically, with the greatest number of Wood Storks being 158 on 30 August 2000 at Red Slough. Three of the five bird censuses we made during the month showed 100 or more storks. On 5 September 2000 we observed 250 Storks, and on 19 September, we counted 350, the highest number ever recorded for Oklahoma. The last date storks were reported at Red Slough in 2000 was on 27 September when we observed eight. All but two reports of Wood Storks in Oklahoma during 2000 were from Red Slough, the other reports being of single birds in Washington and Sequoyah counties in early September (J. Arterburn, pers. comm.).

Red Slough is a 2954-ha impounded shallow fresh marsh project developed in 1998 and 1999. It is located in the Red River floodplain 10 km south of Haworth and 3 km north of the Red River (Fig. 1). The area is enrolled in the U.S. Dept. of Agriculture Wetland Reserve Program, which has resulted in construction of extensive dikes and water level management structures.

A possible factor in the invasion of Wood Storks was the abnormally hot and dry summer in 2000 throughout the south central states, resulting in reduced water levels and concentration of prey species such as fish and crawfish. According to Tom Smith (pers. comm.), the Oklahoma State Forestry fire lookout tower at Tom, 6 km east of Red Slough, recorded only 5.0 cm of rain in July, none in August, and 5.8 cm in late September. Also recorded were 15 days during August when the temperature reached or exceeded 37.8°C. Crawfish, which serve as an important food source




Fig. 1. Location of Red Slough Wetland Reserve Project in McCurtain Co., Oklahoma.

for Wood Storks, are abundant in the shallow impoundments of Red Slough and attract large numbers of wading birds. In mid-August 2000, Heck observed a flock of 98 White Ibis (*Eudocimus albus*) actively feeding on crawfish in 20-cm deep water. The crawfish being captured were approximately 7 cm in length, and approximately 4 seconds would elapse from time of capture of each one until it was swallowed. Heck scanned the flock twice, counting 26 and 28 White Ibis with captured crawfish in their bills, giving an indication of their abundance in the Red Slough impoundments.

The first state record for the Wood Stork in Oklahoma was on 22–26 August 1934, when 1–4 birds were seen near Spiro, LeFlore Co. Until our observations in 2000 the previous high count of Wood Storks for the state was 150 on 13–14 August 1954 in McCurtain Co. (Baumgartner and Baumgartner, Oklahoma bird life, Univ. Oklahoma Press, Norman, 1992). During 1999, we found 27 storks at Red Slough on 31 August and 28 on 13 September, although we did not search extensively for them that year. Storks found in Oklahoma originate from breeding colonies in Mexico (Coulter et al., Wood Stork in The birds of North America (F. Gill and A. Poole, eds.), no. 409, Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union, Washington, D.C., 1999.). Thus, the storks found in Oklahoma probably move through Texas. B. Frenz (pers. comm.) stated that Wood Storks are usually uncommon from mid-June to mid-October and rare at other times in central Texas, but in 2000 many more were present. A raptor migration census during the late summer and fall of 2000 conducted by Hawk Watch International at Hazel Bazemore County Park in northwest Corpus Christi, Texas, included counts of migrating Wood Storks, and in 2000, 875 migrating storks were counted

from 23–27 September.—BERLIN A. HECK, 109 Kaye Dr., Broken Bow, OK. 74728 and W. DAVID ARBOUR, 1462 West Collin Raye Dr., DeQueen, AR. 71832. Received 30 October 2001, accepted 12 November 2001.

*Bull. Okla. Ornithol. Soc.* 34:27, 2001  
© Oklahoma Ornithological Society 2001

**First nesting record for the Common Moorhen in McCurtain County, Oklahoma.**—On 20 July 2000 we observed a family group of two adult and seven flightless young Common Moorhens (*Gallinula chloropus*) in a shallow, flooded marsh in the 2954-ha U.S. Department of Agriculture's Red Slough Wetland Reserve Project in McCurtain County, 10 km south of Haworth, Oklahoma. We found adult Common Moorhens with juveniles in the same area once in August and three times in September. Our last observation of Common Moorhens was on 1 October 2000 when we observed two well-feathered juveniles.

The first documented nesting of Common Moorhens in Oklahoma was in Love County on 15 May 1889 (Nice, *The birds of Oklahoma*, revised ed., Publ. Univ. Oklahoma Biol. Surv. 3:86, 1931). No additional nestings were documented until 1961, when broods were recorded in Bryan and Grady counties. Nestings have since been recorded in Oklahoma, Canadian, Major, Kingfisher, and (now) McCurtain counties (Baumgartner and Baumgartner, *Oklahoma bird life*, Univ. Oklahoma Press, Norman, 1992; J. Arterburn, pers. comm.).—BERLIN A. HECK, 109 Kaye Dr., Broken Bow, OK. 74728 and W. DAVID ARBOUR, 1462 West Collin Raye Dr., DeQueen, AR. 71832. Received 3 November 2000, accepted 31 October 2001.

### Recent Literature

Johnsgard, Paul A., 2001. *Prairie Birds: Fragile Splendor in the Great Plains*. University Press of Kansas. Lawrence, Kansas. ISBN:0-7006-1067-7. 331 pages. \$29.95.

This book celebrates the simple, elegant beauty of the prairies of the American Great Plains and their birdlife. Johnsgard's lifetime of experience, knowledge, and great love of the prairies is quite elegantly expressed in both words and drawings. The book opens with chapters describing the geological structure and plant communities of the different sorts of prairies, information necessary to understand the true complexity of this ecosystem. Johnsgard continues with chapters describing specific bird communities, such as the grassland grouse and prairie-chickens, owls, and prairie sparrows. He ends the book with a discussion of the disappearing prairie and its imperiled avifauna, a sobering account of the current state of conservation in the Great Plains. As always, Johnsgard has illustrated the book with numerous maps and line drawings of birds. These pen-and-ink illustrations are always my favorite part of a Johnsgard book. There is one particularly charming drawing of a Burrowing Owl family and an-

other of an incubating Upland Sandpiper. Throughout the book, Johnsgard has sprinkled quotes from writers such as Peter Matthiessen, William Least Heat-Moon, and Henry David Thoreau. Each chapter ends with a brief section on additional reading, and there is a substantial bibliography and glossary at the end of the book. An appendix describing some of the major grassland preserves in the Great Plains is also included. Johnsgard says it best, "...but a prairie without the song of a Horned Lark or a Sprague's Pipit overhead is no prairie at all. And a place with no prairies at all is not a place that will stir the heart."—*Mary Bomberger Brown*

---

## Index of Bird Names

### Vol. 34, 2001

---

By MARY BOMBERGER BROWN

- Aix sponsa*: 14, 17  
*Archilochus alexandri*: 21  
*Baeolophus griseus*: 12  
    *ridgwayi*: 12  
Bluebird, Eastern: 18  
*Calidris himantopus*: 1  
*Centrocercus minimus*: 11  
    *urophasianus*: 11  
Chickadee, Black-capped: 11  
*Clangula hyemalis*: 11  
*Columbina inca*: 4  
Cowbird, Brown-headed: 8, 18–19  
*Dendrocygna autumnalis*: 13, 17  
Dove, Inca: 4  
Dowitcher, Long-billed: 1  
Duck, Long-tailed: 11  
    Wood: 14, 17  
*Eudocimus albus*: 26  
*Gallinula chloropus*: 27  
*Geothlypis trichas*: 19  
Grouse, Sage: 11  
    Greater Sage: 11  
    Gunnison Sage: 11  
Hummingbird, Allen's: 22  
    Black-chinned: 21, 22  
    Broad-tailed: 21–25  
    Calliope: 22  
    Rufous: 21, 22  
Ibis, White: 26  
*Icterus bullockii*: 12  
    *galbula*: 12, 18  
*Limnodromus scolopaceus*: 1  
*Limnothlypis swainsonii*: 5  
Magpie, Black-billed: 11  
    *Molothrus ater*: 8, 18  
Moorhen, Common: 27  
*Mycteria americana*: 25  
Oldsquaw: 11  
Oriole, Baltimore: 12, 18–19  
    Bullock's: 12  
*Passer domesticus*: 18  
Phalarope, Wilson's: 1  
*Phalaropus tricolor*: 1  
*Philomachus pugnax*: 1  
*Pica hudsonia*: 11  
    *pica*: 11  
*Poecile atricapilla*: 11  
    *atricapillus*: 11  
Reeve: 2  
Ruff: 1–3  
Sandpiper, Buff-breasted: 2  
    Stilt: 1  
*Selasphorus platycercus*: 22  
    *rufus*: 21  
    *sasin*: 22  
*Sialia sialis*: 18  
Sparrow, House: 18  
*Stellula calliope*: 22  
Stork, Wood: 25–27  
Swallow, Tree: 4  
*Tachycineta bicolor*: 4  
Thrush, Wood: 6  
Titmouse, Juniper: 12  
*Tryngites subruficollis*: 2  
Warbler, Swainson's: 5–11  
Whistling-Duck, Black-bellied: 13–17  
Yellowthroat, Common: 19–20